

## Check-List of Additional Taxa to the Supplement Flora of Turkey II

Neriman ÖZHATAY, Şükran KÜLTÜR, Nevriye AKSOY

Faculty of Pharmacy, Department of Pharmaceutical Botany, University of Istanbul, Istanbul-TURKEY

Received: 25.12.1998

Accepted: 05.02.1999

**Abstract:** This checklist contains a total of 168 taxa published as additions to the Flora of Turkey. It includes details of taxa neither recorded by P.H.Davis' "Flora of Turkey and the East Aegean Islands" or its Supplement, nor in the first of this series of checklists. The records in this list are based on data given in nearly 100 publications published up to the end of 1998.

**Key Words:** Turkish flora, new species, additional taxa.

### Türkiye Florasına İlave Edilen Türlerin Listesi II

**Özet:** Türkiye florasına ilave edilen 168 taksonun yeraldiği bir liste hazırlanmıştır. Liste 1998 yılı sonuna kadar yayımlanan yaklaşık 100 çalışma ile P.H. Davis'in Türkiye Florası ve ek cildinde ayrıca Özhata ve arkadaşları tarafından hazırlanan Check-List de yer almayan taksonları içermektedir.

**Anahtar Sözcükler:** Türkiye florası, yeni türler, ilave taksonlar.

### Introduction

The first checklist of taxa additional to the Turkish flora, yet unrecorded in P.H.Davis' "Flora of Turkey and the East Aegean Islands" (1) and its 1988 Supplement (2), was published at the end of 1994 (3). It listed a total of 178 new records/taxa (including 8 new combinations of previously recorded taxa). Since 1994 nearly 100 further papers regarding the flora of Turkey have been published (and in one or two instances traced) and these detail many additional taxa recorded either new to science or as new additions to the Turkish flora. This literature-published in a wide range of journals and other publications – has been drawn together, by the authors and are now housed as a collection in the library of the Department of Pharmaceutical Botany at the University of Istanbul.

This present survey aims to present all the published taxa added to the Turkish flora since the publication of the Flora, its Supplement and the first checklist of additional taxa (the predecessor to this paper). The families and genera are presented in the systematic order presented by the Flora, but species are given in alphabetical order and in an identical format as is given in the Supplement (2). References to the original literature sources are given in bold parentheses at the end of the

taxon name. The list is simply an inventory: the authors have in no way sought to assess the taxonomic status and credibility of new taxa described. Records for new taxa from the East Aegean islands but currently under Greek jurisdiction are not included in the listing.

The list contains a total of 163 taxa, the majority of which are completely new to the flora. This figure comprises 109 taxa new to science (77 species, 26 subspecies, 5 varieties and 1 hybrid) and a further 51 taxa previously unrecorded from Turkey (42 species, 4 subspecies, 4 varieties and 1 hybrid). In addition, 8 taxa previously recorded from the flora have been presented as new combinations and are included here. The following table summarises the statistics of additions to the Turkish flora during the period 1988 to 1998.

We have endeavoured to trace all additional taxa to the flora published before the end of 1998. We recognize that we may have overlooked certain taxa and are fully aware that subsequent to the publications of this paper, further additions will be made to Turkey's exceptionally rich flora. For this reasons, the authors see this checklist as part of a series, with new contributions published every few years. Accordingly we should welcome any additions or comments for inclusion within future issues.

	Check-List I (1988-1994)	Check-List I (1995-1998)	Total (1988-1998)
<b>Taxa new to science</b>			
species	96	77	171
subspecies	16	26	42
variety	11	5	16
hybrid	1	1	2
Total	124	109	233
<b>Taxa new to Turkey</b>			
species	36	42	78
subspecies	8	4	12
variety	2	4	6
hybrid	-	1	1
Total	46	51	97
<b>Previously known taxa</b>			
comb.nov.	3	4	7
stat.nov.	4	3	7
nom.nov.	1	1	2
Total	8	8	16
<b>TOTAL</b>	<b>178</b>	<b>168</b>	<b>346</b>

The list of additional taxa

e = endemic species

\* = new genus for Turkish flora

**PTERIDOPHYTA****19. MARSILEACEAE (1: 62)****\* PILULARIA L. (1:62)****P. globulifera** L., in Sp. Pl. 1100 (1753). [4]

The following information is recorded in the unpublished Flora of Constantinople which is a study by Wladyslaw Jablonowski "1841-1890" [4], as "*Pilularia globulifera* L. from the Asiatic side A2(A) Istanbul in the vicinity of the so-called Sweet Waters of Asia".

**P. minuta** Durieu ex A.Braun, in Monatsber. Koenigl. Akad. (Berlin) 1863: 435 (1864). [5]

(Turkey) W. Anatolia. B1 Izmir: Mares du Mont-Pagus pres Smyrne, v.1866, Balansa 1559 (E).

This specimen is cited by Boissier (1884) in Flora Orientalis [6] but *Pilularia* is not mentioned at all in the Flora of Turkey and the Eastern Aegean Islands [1 & 2]

**AZOLLACEAE**\* **AZOLLA** Lam.

**A. filiculoides** Lam., in Encycl. Meth. Bot. 1: 343 (1783). [7, 8]

European Turkey. A1(E) Kırklareli: Babaeski, Karahalil village, Tekke river, ca. 180-200 m, 22.v.1994, M.Ünal.

This alien species is recorded first time in by Akyar [7] without any locality data, then Ünal & Üzen [8] published with detailed information.

**SPERMATOPHYTA****GYMNOSPERMÆ****21. PINACEAE (1: 67)****4. PINUS L. (1: 72)****P. brutia** Ten., in Prodr. Fl. Nap., 1:57 (1811).

e var.*densifolia*\*, in İ.Ü. Orman Fak. D., Seri A, 39(2): 46 (1989). [9] nom.nud.

Type: (Turkey) C4 Mersin: Gülnar-Pembecik-Karakaya, 400 m. s.n.

*P. nigra* Arnold., in Reise nach Mariazell 8 (1785).  
 subsp. *pallasiana* (Lamb.) Holmboe, in Bergens Mus. Skr. 2, 1 (2): 29 (1914).

e var. *columnaris-pendula*\*, in İ. Ü. Orman Fak. D., Seri A, 39(1): 122 (1989). [10] nom.nud.

Type: (Turkey) C3 Antalya: Alanya Orman İşletmesi, Söğüt İşletme Şefliği, Baloğlu Deresi, Ballı su mevkii, 26.viii.1983, 1400 m. s.n.

\* these new taxa not published as valid

### 23. CUPRESSACEAE (1: 76)

#### 2. JUNIPERUS L. (1: 78)

*J. communis* L., in Sp. Pl. 1040 (1753). subsp. *communis* [11]

European Turkey. A2(E) İstanbul: sur la route de Kalfaköy, ca.1 km de la route principale, 170 m, 7.viii.1969, A.Baytop, ISTE 14902!

### ANGIOSPERMAE

#### DICOTYLEDONES

##### 2. PAEONIACEAE (1: 204)

###### *PAEONIA* L. (1: 204)

*P. mascula* (L.) Miller, in Gard. Dict. No.1 (1768).

e subsp. *bodurii* N.Özhatay, in Karaca Arboretum Magazine, 3(1), 21-22 (1995). [12]

Type: NW Turkey: A1(A) Çanakkale: Çan-Çanakkale, above Çamyayla, 760 m, 29.iv.1992, S.Bodur, ISTE 64073 (holo. ISTE; iso. MARA).

##### 5. PAPAVERACEAE (1: 213)

##### 6. CORYDALIS Medik. (1: 238)

###### *C. caucasica* DC., in Syst. 2: 119 (1821).

e subsp. *abantensis* Liden, in Willdenowia, 26: 26 (1996). [13]

Holotype: (Turkey) Anatolia, A3 Bolu: Bolu-Gerede, 1050 m, beech scrub facing N, flowers pink, 31.iii.1957, Davis & Hedge 26286 (K: isotypus:E).

e *C. lydica* Liden, in Willdenowia, 26: 32 (1996). [13]

Holotype: (Turkey) Anatolia, Boz Dağ, 1700-1800 m, 24.iv.1991, J. & K.Persson 91-96, cultivated in Göteborg Botanic Garden (GB; isotypus: B).

*C. wendelboi* M.Liden, in Notes Roy. Bot. Gard. Edinburgh 45(2): 355 (1988).

e subsp. *congesta* Liden & Zetterlung, in Willdenowia, 26: 26 (1996). [13]

Holotype: (Turkey) Anatolia, A2(A) Bursa: Ulu Dağ, Brian Mathew, s.n. cultivated in Göteborg Botanic Garden (GB).

##### 6. CRUCIFERAEE (1: 248)

##### 24. AETHIONEMA R.Br. (1: 314)

e *A.acarii* Gemici & Leblebici, in Candollea 50(1): 41 (1995). [14]

Type: (Turkey) B2 Kütahya: Gediz, Şaphane Dağı, N part of summit, rocky limestone slopes, 1800-2100 m, 17.vi.1993, Gemici, Y. & G.Görk 7694 (Holo, EGE).

e *A. alanyae* Duman, in Karaca Arboretum Magazine, 2: 172 (1994). [15]

Type: (Turkey) C4 Antalya: Alanya, Gökböl yaylası doğusu, 1600 m, 1.ix.1993, H.Duman and Z.Aytaç 5500 (fl.), (holo-GAZI; iso-ANK, HUB).

e *A. dumanii* M.Vural & N.Adıgüzel, in Doğa Tu Botanik D., 19(4): 481-82 (1995). [16]

Type: (Turkey) B3 Eskişehir: Polatlı to Sivrihisar, 25. km, 870 m, 10.vii.1993, Duman 5011 (holo GAZI; iso ANK).

e *A. karamanicus* Ertuğrul & Beyazoğlu, in Doğa Tu Botanik D., 21(2): 99 (1997). [17]

Type: (Turkey) C4 Karaman: Ayrancı, 2 km W. of Kayaönü (Küçükkoras) Köyü, 1300 m, 6.vii.1988, Ertuğrul 1201 (fl.) (holo. KNYA).

##### 25. THLASPI L. (1: 330)

e *T. leblebicii* Gemici & Görk, in Candollea 50(1): 43 (1995). [14]

Type: (Turkey) Muğla: Köyceğiz, summit of Sandras Dağı, rocky peridotite slopes, 2000-2100 m, 7.vi.1992, Y. Gemici, et al. 6656 (Holo, EGE).

##### 51. ALYSSUM L. (1: 362)

*A. dasycarpum* Stephen ex Willd., in Sp. Pl. 3(1): 469 (1800).

var. *minus* Bornm. ex Dudley, in Notes Roy. Bot. Gard. Edinb., 23(2): 157, fig. 1A (1962). [18]

W. Anatolia. B3 Eskişehir: Seyitgazi, Aslanbeyli, stony slopes, 1050 m alt., 10.v.1992, Orcan 6846 (AUF).

*A. minus* (L.) Rothm., in Feddes. Rep. 50: 77 (1941).

var. *collinum* (Brot.) Townsend, in Taxon 16: 467 (1967). [18]

W. Anatolia. A3 Eskişehir: Sarıcakaya, Dağküplü-Mayıslar, 1 km to Dağküplü, fields, 550 m alt.,

22.v.1990, Orcan 4961 (AUF); *ibid.*, Orcan 4962 (AUF); B1 Balıkesir: Burhaniye, roadside, 50 m alt., 20.iv.1991, Orcan 5813 (AUF), B3 Eskişehir: Eskişehir to Sarıcakaya, 2 km to Yarımca village, roadside, 1100 m alt., 19.v.1991, Orcan 5947 (AUF); *ibid.*, Orcan 5950 (AUF).

73. *ERYSIMUM* L. (1: 466)

E. *rhodium* Snogerup, in Opera Bot. 13: 41 (1967). [19]

SW. Anatolia: The Marmaris Peninsula.

\* *LUNARIA* L.

*L. annua* L., in Sp. Pl. 653 (1753).

subsp. *annua* [20]

European Turkey. Al(E) Kırklareli: Karataş mahallesi, pastures, 250 m, 25.iv.1990, Ş. Yıldırımlı 13445 F. Nayır (HUB; Herb. Yıldırımlı).

Alien species.

10. *VIOLACEAE* (1: 524)

*VIOLA* L. (1: 524)

V. *bocquetiana* Yıldırımlı, in Ot Sistematisk Botanik D., 1(1): 2 (1994). [21]

Type: (Turkey) B7 Tunceli: Ovacık, Karagöl valley, rock crevices in stony slopes, 1800 m, 30.vi.1983, Ş. Yıldırımlı 5418 (Holo-HUB; iso. Herb. Yıldırımlı).

13. *CARYOPHYLLACEAE* (2: 15)

2. *MINUARTIA* L. (2:38)

M. *asiyeae* Duman, in The Karaca Arboretum Magazine, 4(4): 159 (1998). [22]

Type: Turkey C4 Antalya: Alanya, Elmalısu-Gökbeş yaylası, limestone rocks, 1250 m, 18.vii.1995, H.Duman, K.H.C.Başer & A. Altıntaş, HD 5836 (holo.GAZI, iso.E. ESSE, HUB, AIBU).

18. *DIANTHUS* L. (2: 99)

e D. *akdagensis* Gemici & Leblebici, in Candollea 50(1): 43 (1995). [14]

Type: (Turkey) C2 Muğla: Fethiye, Akdağ, W side, above Camialanı, 2200 m, rocky limestone slopes, 8.vii.1993, Y. Gemici, G.Görk & N.Özel 6928 (Holo. EGE).

D. *elegans* d'Urv., in Enum. Pl. Ins. Pont. Eux., 46 (1822).

e var. *gramineus* R.M.Burton, in The Karaca Arboretum Magazine, 3(4): 184-185 (1997). [23]

Holotype: (Turkey) C2 Muğla: Asar, shady cliff above north side of Ködürümüşü Limanı, 14 km south of Fethiye,

28.v.1991, R.M.Burton An. 206 (Leafy branch with dead flowering shoot of previous year) (AIBU).

24. *GYPSOPHILA* L. (2: 149)

G. *glomerata* Pallas ex Bieb., in Fl. Taur.-Cauc. 1: 321 (1808). [24]

European Turkey. A1(E) Edirne: 1 km from Süloğlu to Lalapaşa, 7.viii.1975, N. and E.Özhatay, ISTE 33122.

29. *SILENE* L. (2: 179)

S. *caryophylloides* (Poir.) Otth., in DC., Prodr. 1: 369 (1824).

e subsp. *binbogaense* Duman et Vural, in Doğa Tu Botanik D., 19(4): 478 (1995). [25]

Type: (Turkey) B6 Kahramanmaraş: Göksun, Doğankonak köyü, Binboğa Dağı, Kapıkaya batı sırtları, 1800-2150 m, step, 2.vii.1992, Duman et Aytaç 4567 (holo-GAZI, iso-ANK, HUB).

e S. *denizliense* Aytaç, in Thaiszia, J.Bot., Kosice, 8: 7-11 (1998). [26]

Type: Turkey. C2 Denizli: Çamlık, around taşocağı, 830 m, 6.vi.1995, on a calcareous rocky site (Aytaç 6927), (holo. GAZI, iso. ANK, HUB). Para.: *ibid.* Aytaç 6764 & Karavelioğulları, 27.ix.1994.

15. *POLYGONACEAE* (2: 265)

6. *POLYGONUM* L. (2: 269)

e P. *ekimianum* Leblebici, H.Duman & Aytaç, in Willdenowia 23: 163-165 (1993). [27]

Type: (Turkey) C6 Kahramanmaraş: Ahır Dağı, Karagöl mevkii, step, 1650 m, 19.x.1990, Duman et Aytaç 3411 (Holo-EGE; iso-ANK, B, GAZI, HUB).

7. *RUMEX* L. (2: 281)

R. *arifolius* All., in Melang. Phlos. Math. Soc. Roy. Turin (Misc. Taur) 5: 94 (1774). [28]

N. E. Turkey. A6 Sivas: Şerefiye, Köse Dağı, Cehennemdere, su kenarı ve çayırlıklar, 1800 m, 11.viii.1984, B.Yıldız 5005, N.Çelik. A7 Sivas: Suşehri, Aksu köyü, Köse Dağı, yaprak döken orman, 1700-1800 m, 26.vi.1985, B.Yıldız 5703.

16. *CHENOPodiaceae* (2: 294)

8. *CERATOCARPUS* L. (2: 313)

C. *turkestanicus* Sav.-Rycz., in Addenda, V. p.874. [29]

E. Turkey. B10 Ağrı: Doğubeyazıt, Gürbulak, Iran frontier, steppes, 1550 m, 21.vii.1990, Ş. Yıldırımlı 13752 (HUB, Hb. Yıldırımlı); *ibid.*, 20.vi.1991, Ş.Yıldırımlı 14538 (HUB; Hb. Yıldırımlı).

22. *SALSOLA* L. (2: 328)
14. *S. canescens* (Moq.) Spach, in Fl. Or. 4: 963 (1879).
- e subsp. *serpentinicola* Freitag & E.Özhatay, in Willdenowia, 27: 185-187 (1997). [30]
- Holotype: Turkey, C2 Muğla: distr. Köyceğiz, Sandras Dağı, Serçe Gediği, 1650 m, open *Pinus nigra* forest, serpentine rock, 6.ix.1978, N. & E.Özhatay (ISTE 51122; iso. ISTO 20668, KAS).
22. *TAMARICACEAE* (2: 348)
2. *TAMARIX* L. (2: 349)
- T. tetragyna* Ehrenb., in Linnaea, 2: 247-258 (1827). Syn: *T. brachystachys* Bunge, *T. meyeri* Boiss. [31]
- Southern Turkey.
27. *GUTTIFERAE* (2: 355)
1. *HYPERICUM* L. (2:355)
- e *H. kazdagensis* Gemici & Leblebici, in Candollea, 50(1): 46 (1995). [14]
- Type: (Turkey) B1 Balıkesir: Edremit, Kazdağı, NE of Sarıkız Tepe, scree, 1500 m, 23.ix.1993, Y. Gemici, et al. 15259 (Holo, EGE).
- e *H. peshmenii* Yıldırımlı, in Ot Sistematič Botanik D., 2(2): 12 (1995). [32]
- Type: Türkiye. B7 Erzincan: Kemaliye, Başpınar, Armağan Köyü, Karlık çeşme, around Toptaş, crevices of limestone rock, 1750 m, 21.viii.1980, Ş.Yıldırımlı 3902 (Holo. HUB, iso. Herb. Yıldırımlı).
30. *LINACEAE* (2: 425)
1. *LINUM* L. (2: 425)
- L. maritimum* L., in Sp. Pl. 280 (1753). [33]
- S.W. Anatolia. C2 Muğla: Köyceğiz, Hamitköy, 10 m, 17.vii.1991, A. Güner 9626.
31. *GERANIACEAE* (2: 450)
3. *ERODIUM* L'Hérit. (2: 475)
- e *E. birandianum* İlarslan & Yurdakulol, in Flora Mediterranea 4: 167 (1994). [34]
- Type: Turkey (A5 Kastamonu): Devrekani, Yaralığöz Mountain, subalpine limestone rocks, 1900 m, 9.vii.1991, E.Yurdakulol 3552 (holo. HUB, GAZI; iso. ANK).
- e *E. hendrikii* K.Alpınar, in Edinb. Jour. Bot., 51(1): 68-69 (1994). [5]
- Type: (Turkey) A7 Gümüşhane: Yağmurdere, 1800 m, 10.viii.1989, K.Alpınar et H.t Hart, ISTE 61049 (holo. ISTE; iso.E).
- e *E. olympicum* Gemici & Leblebici, in Candollea, 50(1): 46 (1995). [14]
- Type: A2 Bursa: Uludağ, N part of Zirve Tepe, limestone cliff crevices, 2300 m, 5.viii.1993, Y. Gemici, et al. 8280-b (Holo, EGE).
45. *LEGUMINOSAE* (3: 1)
11. *CHAMAECYTISUS* Link (3: 16)
- e *C. gueneri* Duman, Başer & Malyer, in Karaca Arboretum Magazine, 4(1): 13-15 (1997). [35]
- Type: Turkey. C2 Muğla: Köyceğiz, Sandras Dağı, Ağla-Eskere, 1700 m, 25.vii.1992, open *Pinus nigra* forest, serpentine slopes, A. Güner 10760-A.A. Dönmez & H. Şağban (holo. HUB, iso. GAZI, E).
- C. wulfii* (V.Krecz.) A. Klaskova, in Preslia 30: 214 (1958). [36]
- C. Anatolia. A4 Ankara: Kızılcahamam; Çamçukuru Orman evi, *Pinus nigra* ormanı, 8.v.1966, A. Baytop (ISTE 9211); A4 Kastamonu: Ilgaz Da., 1700-750 m, 2.vii.1989, M.Koyuncu 15206 (AEF); A4 Kastamonu, Ilgaz Da., *P. nigra* ormanı, 1465 m, 19.v.1972, A. ve T. Baytop (ISTE 21726); A5 Kastamonu: Tosya, Dedem Da. around Dereşeki, *P. nigra* ormanı, ca.1450 m, 31.vi.1976, M. Kılıç 6264 (ANK); B2 Kütahya: Murat Da., Kesik Söğüt Yaylası (Kızılıluk-Kurugöl), stony places, 1800-1900 m, 3.vii.1976, A. Baytop, K. Alpynar (ISTE 35065); B2 Kütahya: Yellice Da., stony places, 1550 m, 5.vi.1994, H. Malyer 9324 (BULU) and 7236 (ESSE); B3 Afyon: Bayat, north side of Köreoğlu Tepesi, ca.1300 m, 24.v.1975, M. Vural, 137 (ANK).
32. *ASTRAGALUS* L. (3: 49)
- e *A.akmanii* Aytaç et Duman, in Doğa Tu Botanik D., 19(4): 477-478 (1995). [25]
- Type: (Turkey) C6 Kahramanmaraş: Ahır Dağı, Karagöl mevkii, 1750-1800 m, 31.vii.1991, Aytaç et Duman 4219 (holo-GAZI, iso-ANK, HUB).
- e *A. beypazaricus* Podlech & Aytaç, in Sendtnera, 5: 265-268 (1998). [37]
- Holotype: Turkey. A4 Ankara: 12 km from Beypazarı to Nallıhan, 600 m, 2.vii.1997, Aytaç 7610 (GAZI; Iso: MSB).
- e *A. nigrifructus* Podlech & Aytaç, in Sendtnera, 5: 265-268 (1998). [37]

Holotype: Turkey. C4 Konya: Ereğli, İvriz, 1200-1300 m, 22.vi.1997, Aytaç 7600 (GAZI !; Iso: MSB !).

e *A. serpentinicola* Duman & Ekim, in Willdenowia, 27: 181-183 (1997). [38]

Type: (Turkey) C2 Burdur, Yeşilova, S of Salda lake, clearings in *Pinus nigra* and *Quercus* forest, on serpentine, 1170-1200 m, 11.vii.1993, H.Duman 5078 & F.A. Karavelioğulları (GAZI; iso. ANK, B, ISTE).

41. *LENS* Miller (3: 325)

e *L. tomentosus* Ladizinsky, in Bot. Journal of Linnean Society, 123: 260 (1997). [39]

Type: (Turkey) C8 Mardin: 22 km north-west of Mardin, collected by G. Ladizinsky, accession No. 133, grows in Rehovot, Israel, Holo. HUJ, iso.E.

42. *LATHYRUS* L. (3: 328)

*L. inconspicuus* L., in Sp. Pl. 730 (1753).

var. *stenophyllus* (Boiss.) Rech. f., in Fl. Iranica 74 (1979). [40]

S.E. Anatolia. D8 Diyarbakır, Lice, Lice'den Kulp'a doğru 5 km, 850 m, 8.vi.1985, A.S. Ertekin, DUF 4730, Lice, Şeyhan köyü çevresi, 910 m, 8.v.1991, A.S. Ertekin, DUF 6847. C7 Ş.Urfâ, Siverek, Bahser köyü, 900 m, 31.v.1986, A.S. Ertekin, DUF 5163, Siverek, TRT vericisinden Karabahçe'ye doğru 11 km, Karacadağ, 1500 m, 7.v.1981, Ö. Saya, DUF 2457, 15.v.1984, A.S. Ertekin, DUF 4001, Siverek, TRT vericiyi yol ayırmayı çevresi, Karacadağ, 1260 m, 15.v.1984, A.S. Ertekin, DUF 3988, Siverek, Siverek'ten Diyarbakır'a doğru 13 km, 1060 m, 3.vi.1988, A.S. Ertekin, DUF 5322, Diyarbakır, Ş.Urfâ'ya doğru 39 km, Karaca Dağı'nın kuzey etekleri, 1100 m, 15.v.1984, A.S. Ertekin, DUF 3974, Çınar, Hatuni köyü çevresi, Karaca Dağı'nın doğu etekleri, 780 m, 19.v.1989, A.S. Ertekin, DUF 5882.

46. *TRIFOLIUM* L. (3: 384)

*T. ornithopodioides* (L.) Sm., in Fl. Brit. 2: 782 (1800). Syn: *Melilotus ornithopodioides* L., in Sp. Pl. 768 (1753). *Trigonella ornithopodioides* Desr., in Lam. Encycl. Meth. Bot. 4: 67 (1797). [41]

European Turkey. A1(E) Edirne: Keşan, SW of Mecidiye, at the eastern end of the beach extending east of lake Tuzla, 5 m, 19.iv.1994, A.J. Byfield & N. Özhatay (AJB 326), det. A.J. Byfield, ISTE 66433.

50. *MEDICAGO* L. (3: 483)

*M. heyniana* Greuter, in Candollea 25: 190 (1970). [19]

SW. Anatolia: The Marmaris Peninsula.

46. *ROSACEAE* (4:1)

21. *ALCHEMILLA* L. (4: 80)

e *A. ancerensis* Kalheber, in Sendtnera, 2: 412 (1994). [42]

Type: Turcia, prov. A8 Rize: İkizdere, Ballıköy (Anzer), 2150 m, pratum, 19.viii.1984, leg. M. Vural 3090. Herb. Univ. Hacetepensis, Ankara (holo, HUB).

e *A. cimilensis* Kalheber, in Sendtnera, 2: 404 (1994). [42]

Type: Turcia, prov. A8 Rize: İkizdere, Cimil, Basköy üstleri, 2100 m, pratum, 24.vii.1984, M. Vural 3220 B (HUB, holo).

*A. daghestanica* Juz., in Delect. Semin. Hort. Bot. Acad. Sc. URSS (1934) 5. [43]

N.E. Anatolia. A7 Trabzon: East slopes of Zigana Dağ Tesisleri (together with some species of *Pinus*, *Rhododendron ponticum*, *R.luteum* and *Delphinium*), 1750-1800 m, 10.vii.1994, S.Hayırlıoğlu 81, KTÜB.

e *A. elevitensis* Kalheber, in Sendtnera, 2: 406-407 (1994). [42]

Type: Turcia, prov. A8 Rize: Çamlıhemşin, Elevit, Cevizli vallis, ripa rupester rivi, 2100 m, 15.vii.1985, M. Vural 3512 (holo, HUB).

*A. epipsila* Juz., in Grossh., Fl. Kavk. IV (1934) 328. [43]

N.E. Anatolia. A8 Trabzon: on the road of Sürmene to Arpalı Köyü near stream (together with some species of *Carex*, *Achillea*, *Bromus*, *Secale*, among others), 2000 m, 1.viii.1994, S.Hayırlıoğlu 41, KTÜB.

e *A. hemisinica* Kalheber, in Sendtnera, 2: 408-409 (1994). [42]

Type: Turcia, prov. A8 Rize: Çamlıhemşin, Ortaköy, Vercenik Dagı, Alpinik küçük dere yatağı, 2700 m (tüylü), 13.viii.1985, M. Vural 3685 (holo, HUB).

*A. holotricha* Juz., in Addenda, Fl. USSR 10: 295 (1941). [43]

N.E. Anatolia. A7 Trabzon: East slopes of Zigana Dağ Tesisleri (together with some species of *Euphorbia*, *Rhododendron luteum* and *Pinus sylvestris*), 1750-1800 m, 10.vii.1994, S.Hayırlıoğlu 84, KTÜB.

e *A. ikizdereensis* Kalheber, in Sendtnera, 2: 405-406 (1994). [42]

Type: Turcia, prov. A8 Rize: İkizdere, Dereköy-Yerelma vicus, inclinatio vallis, pratum, 1000 m, 10.vi.1984, M.Vural 3008 (1) (holo, HUB).

e *A. kackarensis* Kalheber, in Sendtnera, 2: 407-408 (1994). [42]

Type: Turcia, prov. A8 Rize: Çamlıhemşin, Yukarı Kavrun, Kaçkar Dağı, Narduu stricta caespes, 2350 m, 7.viii.1985, M. Vural 3636 (holo, HUB).

*A. smirnovii* Juz., in Addenda, Fl. USSR 10: 295 (1941). [43]

N.E. Anatolia. A7 Trabzon: Along to Değirmendere valley above Zigana Dağ Tesisleri (together with some species of *Euphorbia*, *Rhododendron luteum* and *Pinus sylvestris*), 1750 m, 10.vii.1994, S. Hayırlıoğlu 110, KTÜB.

*A. speciosa* Buser apud Magnier, in Scrinia fl. Selectae (1893) 281. [43]

N.E. Anatolia. A7 Trabzon: Zigana Geçidi and surroundings together with some species of *Euphorbia*, *Malva sylvestris*, among others, 1800 m, 17.vii.1994, S. Hayırlıoğlu 87, KTÜB.

e *A. trabzonica* Hayırlıoğlu-Ayaz & Beyazoğlu, in Ann. Bot. Fennici, 34: 111-112 (1997). [44]

Holotype: (Turkey) A7 Trabzon: Zigana Pass, S and SW meadows, by streamside banks (together with some species of *Rhododendron*, among others), 1750 m, 20.vi.1995, S. Hayırlıoğlu (Ayaz) 190 (KTÜB).

#### 34. PYRUS L. (4: 160)

e *P. serikensis* A.Güner & H.Duman, in Karaca Arboretum Magazine, 2: 166 (1994). Syn.: *Pyrus boissieriana* Buhse subsp. *crenulata* Browicz, in Notes Roy. Bot. Gard. Edinb., 31: 322 (1972). [45] stat. et nom. nov.

Type: (Turkey) C3 Antalya: 17 km S. of Serik, c. fr., 16.viii.1959, M. and D. Zohary 3092 (holo.HUJ).

SW. Anatolia. (Turkey) C3 Antalya: 17 km S. of Serik, 16.viii.1959, M. and Zohary 3088, 3091. Aksu, Ortaköy çevresi, 100-150 m, 28.viii.1993, H. Duman 5411 (fr.) ! Serik, Dikmen Köyü, 10 m, 19.iii.1994, H. Duman 5556! *ibid.*, 24.x.1993, A. Güner 11641 (fr.)! Aksu, Serik arası, Aşağıkocayatak Köyü yol ayrımı, 20 m, 19.iii.1994, H. Duman 5558 !

#### 49. LYTHRACEAE (4: 174)

##### 1. LYTHRUM L. (4: 174)

e *L. anatolicum* Leblebici & Seçmen, in Doğa Tu Botanik D., 19(5): 559-560 (1995). [46]

Type: (Turkey) A3 Bolu: Düzce, S. of Efteni (Melen) Gölü, aquatic marsh. ca.140 m, 27.vii.1985, E. Leblebici 6032, Ö. Seçmen (EGE 5102).

#### 50. ONAGRACEAE (4: 180)

##### 3. LUDWIGIA L. (4: 182)

*L. peploides* (Kunth) P.H. Raven, in Reinwardtia, 6: 393 (1963). [47]

South Turkey. C3 Antalya: Antalya-Kemer road, 10 km, Sarısu forest Camp, in rivers, 5 m, 19.vii.1997, R.S. Göktürk 4226. C3 Antalya: Sarısu, 5 m, H. Peşmen 4916.

Alien species.

#### 55. CUCURBITACEAE (4:200)

##### \* SICYOS L.

*S. angulatus* L., in Sp. Pl. 1013 (1753). [48]

N.E. Anatolia. A8 Artvin: Borçka, Maçahel (Camili) çevresi, bahçe kenarı, 370 m, 21.viii.1994, A. Güner 11700, H. Duman, F. Karavelioğulları, R. Disperati

Alien species.

#### 58. CRASSULACEAE (4: 209)

##### 4. SEDUM L. (4: 224)

e *S. aytacianum* J.Metzger, in Cactus & Succulent Journal (V.S), 66: 259-260 (1994). (Sect. *Sedum* L.; Sect. *Eurasistica* Kyphocarpia Fröderström). [49]

Type: A7 Trabzon: from limestone rock on a mountain-range ca.3 km west of Maçka, 1800-1850 m, 21.v.1992, J. Metzger 2046.1 (holo GAZI, cult.J. Metzger from J. Metzger 2046.1; iso BASBG and Metzger herbarium, Basle. Cult. GAZI; Basle Botanic Garden; Vaclav Plestil in Turnow and Ladislav Hlavaty in Prerov, Czech Republic.

*S. eriocarpum* Sibthorp & Smith, in Fl. Graec. 1, 310 (1809).

e subsp. *caricum* (Carlström) 't Hart, Ot Sistematisk Botanik D., 2(2): 8 (1995). [50] com. nov.

Typus: Turkey, prov. C2 Muğla: Armutalanı, c.3 km W of Marmaris, open ground on serpentine, Carlström 9694, 23 Apr. 1983 (LD).

e *S. vuralianum* J.Metzg., in British Cactus & Succulent Society Journal, 11(4): 100-101 (1993). (Sect. *Sedum*) [51]

Type: A7 Trabzon: Zigana in open ascending terrain with limestone boulders strewn in, with some trees of *Pinus sylvestris*, 200 m, E of the N tunnel entrance, 2050 to 2070 m, 23.v.1992, J. Metzger 2012.1 (holo GAZI, cult. J. Metzger from J. Metzger 2021.1), Hb. Metzger, Basle; cult. GAZI, Basle Botanic Garden & Tumov, Czech Republic.

59. SAXIFRAGACEAE (4: 249)

1. SAXIFRAGA L. (4: 249)

S. graeca Boiss. et Heldr., in Boiss., Fl. Or. 2: 807 (1872). [52]

European Turkey. A1(E) Edirne: 2 km from Lalapaşa, Bağlık deresi, 25.iv.1974, N. and E. Özhatay, ISTE 27611, same place 22.v.1978, N. and E. Özhatay, ISTE 39506 (rev. V.A. Matthews); Kırklareli: Kırklareli to Dereköy, Koruköy, 400 m, 22.v.1978, N. and E. Özhatay, ISTE 39530.

This species (*S.graeca* Boiss.) is recorded in P.H.Davis "Flora of Turkey and East Aegean Island" vol.4, p.254 from East Aegean Islands (Samos, Ikaria, Khios).

63. UMBELLIFERAE (4: 265)

I. HYDROCOTYLE L. (4: 288)

H. vulgaris L., in Sp. Pl. 234 (1753). [33]

S.W. Anatolia. C2 Muğla: Köyceğiz, Lambataklığı çevresi, 15 m, 20.v.1991, A. Güner 9137.

7. ECHINOPHORA L. (4: 306)

e E. lamondiana B. Yıldız & Z. Bahçecioğlu, in The Karaca Arboretum Magazine, 4: 7-8 (1997). (Sect. Chrysophora DC.) [53]

Type: Turkey B6 Malatya: 4-5 km from Hemikhan to Hasançelebi, eroded slopes, 1100 m, 26.ix.1992, B.Yıldız 9872 (holo GAZI, iso. ISTE).

31. SESELI L. (4: 367)

S. gummiferum Pallas ex Smith, in Sp. Pl. 251 (1753).

subsp. *gummiferum* [54]

C. Anatolia. A4 Ankara: İdris Dağı, Adilahmet köyü üstü, 1700 m, 9.ix.1988, M.Koyuncu (AEF 16347); Hasanoğlan, İdris Dağı, Hasanoğlan'ın üstü, 1500-1600 m, 27.vi.1989, Koyuncu et al. (AEF 16348).

42. PRANGOS Lindl. (4: 382)

P. longistylis (Boiss.) Pimenov & Kluykov, in Candollea 52(2): 110 (1997). Syn.: *Ferulago longistylis* Boiss., in Ann. Sci. Nat. Bot. Ser. 3, 1: 321 (1844), *Prangos platychaena* Boiss. ex Tchich., Asie Min., Bot. 1: 457 (1860). [55] com. nov.

Holotype: B8 Erzurum: "Armeria: prope pagum Kale, haud procul am. Mille Lacuum (Bingoeldagh), alt. 1900-2000 m, 1858, Tchihatcheff 896 (P).

80. TORDYLIUM L. (4: 504)

e T. ebracteatum Al-Eisawii & Jury, in Bot Jour. of

Linnean Society, 97: 401-402 (1988). [56]

Type: (Turkey) Muğla, limestone sea-cliffs just before entrance of Bozburun Harbour, 50-100 m, 8.vi.1962, D. Dudley 35477 (E).

67. VALERIANACEAE (4: 550)

1. VALERIANA L. (4: 551)

V. montana L., in Sp. Pl. 32 (1753). [57]

N.E. Anatolia. A8 Rize: Çamlıhemşin, Yukarı Kavrun yayası, Derebaşı mevkii, çayırlık, 2750 m, 5.viii.1985, A. Güner-M. Vural (A. Güner 6705) (AEF 11895)

69. DIPSACACEAE (4: 582)

2. CEPHALARIA Schrader ex Roemer & Schultes (4: 585)

e C. ekimiana R.S. Göktürk & H. Sümbül, in Ann. Bot. Fennici 34: 153 (1997). [58]

Holotype: (Turkey) S. Anatolia, C4 Karaman, Ermeneke, Tepebaşı village, roadside, on disturbed ground, 1100 m, 2.viii.1995, R.S.Göktürk 3479 (Akdeniz University Herbarium; isotypes HUB, ANK, GAZI). Paratype: (Turkey) S. Anatolia C4 Karaman: Ermeneke, Tepebaşı village, roadside, on disturbed ground, 1100 m, 6.viii.1996, R.S. Göktürk 3892 (Akdeniz University Herbarium).

70. COMPOSITAE (5: 1)

5. AMBROSIA L. (5: 47)

A. elatior L., in Sp. Pl. 987 (1753). [59]

N.E. Anatolia. A7 Trabzon: 1 km W. of Çarşıbaşı, damp roadside grassland, 10-20 m, 23.viii.1995, A.J. Byfield and S.Atay (ISTE 69357).

Alien species.

16. HELICHRYSUM Gaertner (5: 80)

H. plicatum DC., in Prodr. 6: 183 (1838).

e subsp. *isauricum* Parolly, in Willdenowia, 25: 239-241 (1995). [60]

Type: (Turkey) C4 Karaman, Oyüklu Dağ bei Ermeneke, Nordflanke, 36°51'N, 32°53'E, subalpine Grasflur mit *Senecio cilicicus*, 1900 m, feinerdreich, Ophiolitikh, 25-30, SE-exp., 11.vii.1992, Parolly 92-45 b-1 (BSB, iso-B).

20. LOGFIA Cass (5: 107)

L. minima (Sm.) Dumort, in Fl. Belg. 68 (1827). [61]

European Turkey. A1 (E) Kırklareli: Kırklareli to Kofçaz, at the turning to Kadıköy, edge of a *Helianthus annuus* field, 3.vii.1975, A. Baytop, ISTE 32501. A1(E)

Kırklareli: SW of İğneada, Longoz forest, dry acid grassland, on river terrace gravels, 10 m, vi.1993, A.J.Byfield B 2354, ISTE 69708. A1(E) Edirne: Enez, 4 km S of Küçük Evren toward sea, sandy tracks through young *Pinus* plantation, 20 m, 1.vi.1994, A.J.B yfield B 653, ISTE 66762. A1(E) Edirne: Enez to Ipsala, Karpuzlu, E of Sığırıcı Lake, short grazed sandy pasture, 30 m, 1.v.1994, A.J. Byfield B 662, ISTE 66771. C2 Denizli: 5 km above Babadağ, N dry shale slopes, light acid soils, 1300 m, 27.v.1995, Iter Anatolicum 95/054, A.J. Byfield, s.n., ISTE 70344.

#### 42. ANTHEMIS L. (5: 174)

A. melanolepis Boiss., in Fl. Or. Suppl. 297 (1888). [33]

S.W. Anatolia. C2 Muğla: Köyceğiz, Yangı Köyü sırtları, 70 m, 20.iv.1991, A. Güner 8947.

#### 60. ONOPORDUM L. (5: 356)

O. majori Beauverd, in Bull. Soc. Bot. Genève ser. 2, 6: 152 (1914). [33]

S.W. Anatolia. C2 Muğla: Köyceğiz, Ekincik, Sandallı tepe, 700 m, 16.vii.1991, A. Güner 9598. Dalyan, Eskiköy, 20 m, 15.vii.1991, A. Güner 9634.

O. sirsangense Rech. f., in Fl. Iranica, 139 a: 164 (1979). [62]

E. Anatolia. C9 Van: 15 km from Çatak to Narlı, river banks, 1450 m, 14.viii.1984, E.Tuzlaci, ISTE 54964!

#### 62. CIRSIUM Miller (5: 370)

e C. dirmilense R.M. Burton, in Karaca Arboretum Magazine, 3(3): 123-124 (1996). [63]

Holotype: (Turkey) C2 Burdur: Altinyayla side of Dirmil Geçidi, c. 1500 m, serpentine, 23.x.1993, R.M.Burton An. 902 (ABANT).

#### 115. HIERACIUM L. (5: 696)

e H. nydeggerorum Gottschlich, in Bauhinia, 11/4: 243-244 (1996). [64]

Holotype: (Turkey) A9 Erzurum: Olur, Karadağ, Föhrenwald, 1700 m, 21.vii.1984 , M. Nydegger 19343 (G).

#### 118. PRENANTHES L. (5:763)

e P. oyukludaghensis Parolly, in Willdenowia, 25: 242-244 (1995). [60]

Type: (Turkey) C4 Karaman, Oyüklu Dağ bei Ermenek, Nordostflanke, Kar, 36°51' N, 32°53' E, subalpine steinschuttflur mit *Scrophularia myriophylla* und *Ferulago trachycarpa*, 2000 m, feinerdereicher,

ruhender Mittelschutt, Dolomit, 30, SE-exp., 23.vii.1992, Parolly A354-1 (BSB, iso-B, herb. Parolly).

#### 72. CAMPANULACEAE (6: 2)

##### 1. CAMPANULA L. (6: 2)

e C. damboldtiana Davis & Sorger, in Notes RBG, 37(2): 265 (1979). (Sect. Dictyocalyx (Fed.) Damboldt). [65]

Type: (Turkey) A4 Ankara: 16 km E. Ayaş, sandiger Steppenhang, 1190 m, 7.vii.1977, F. Sorger 77-56-9 (holo. Hb. Sorger; iso. E).

C. latiloba A. DC., in DC., Prodr., 7: 478 (1839).

e subsp. rizeensis A. Güner, in Candollea, 39: 348 (1984). [66]

Type: (Turkey) A8 Rize: Çamlıhemşin çevresi, sarp kayalık, c.250 m, 21.v.1981, A. Güner 3547 (Holotype HUB; isotype ANK).

e C. kirikkaleensis A.A. Dönmez & A. Güner, in Karaca Arboretum Magazine, 2: 115-118 (1993). (Sect. Rupestres (Boiss.) Charadze) [67]

Type: (Turkey) B5 Kırıkkale: Delice çevresi, cipsli yamaçlar, bozkır, 720 m, 17.viii.1993, A.A. Dönmez 3924, M. Açıkgöz, C. Uyanık (holo. HUB, iso.GAZI, ANK.).

C. rotundifolia L., in Sp. Pl. 163 (1753). [5]

European Turkey. A1(E) Kırklareli: 2 km from Demircihalil Village to Dereköy, 200 m, 21.viii.1988, K.Alpınar & H.'t Hart (ISTE 59685).

#### 73. ERICACEAE (6: 89)

##### 1. RHODODENDRON L. (6: 90)

R. ponticum L., in Sp. Pl. ed. 2:562 (1762). subsp. ponticum

e var. heterophyllum Anşın, in Tr. J. of Agricultural and Forestry, 18: 139 (1994). [68]

Holotype: (Turkey) A8 Trabzon: Trabzon-Of, Kumludere Köyü, 50-100 m, *Quercus-Carpinus* yapraklı ormanlarında, *Laurocerasus*, *Rhododendron* ve *Corylus* çalılıkları ile birlikte bulunmaktadır. S. Terzioğlu, 10.xi.1992 (KATO:9884).

#### 87. BORAGINACEAE (6: 237)

##### 18. BUGLOSSOIDES Moench (6: 315)

B. glandulosum (Velen) R.Fernandez, in Bot.J.Linn. Soc., 64: 379 (1971). [69]

Type: (Turkey) A4 Ankara: Kızılcahamam, Çamkoru, Kadınboğazı, moist meadows, 1400-1450 m, 8.v.1986, Ş. Yıldırımlı, 9193 (HUB, Herb. Yıldırımlı).

20. ECHIUM L. (6: 318)  
E. arenarium Guss., in Ind. Sem. Horto Boccad. 5 (1826). [19]  
S.W. Anatolia: The Marmaris Peninsula
88. SOLANACEAE (6: 437)  
4. PHYSALIS L. (6: 444)  
P. pubescens L., in Sp. L. 183 (1753). [59]  
N.E. Anatolia. A8 Artvin: Kemalpaşa, beach W. of river mouth, sandy dry grassland behind beach, 5 m, 17.viii.1995, A.J. Byfield and S. Atay (ISTE 69309).  
Alien species.
89. SCROPHULARIACEAE (6: 458)  
17. VERONICA L. (6: 689)  
V. peregrina L., in Spec. Pl. 14 (1753). [70]  
S.E. Anatolia. C5 Adana: Seyhan Dam Lake, Karaömerli village, shorelines, wet places, 60-70 m, 16.xii.1990, H. Çakan 2338 (ADA).
94. LABIATAE (7: 36)  
2. TEUCRIUM L. (7: 53)  
e T. ekimii Duman, (Sect. Teucrium Bentham), in The Karaca Arboretum Magazine, vol. 4(3): 125-130 (1998). [71]  
Type: Turkey. C2 Antalya: between Beldibi and Kemer, Kuruçay Dere, 40-50 m, shady limestone cliffs, 10.vi.1996, H. Duman 6423, N. & E. Özhataş (holo. GAZI; iso. E, ISTE, ANK).
16. BALLOTA L. (7: 156)  
e B. sechmenii Gemici & Leblebici, in Candollea, 50(1): 50 (1995). [14]  
Type: (Turkey) C4 Antalya: N part of Anamur, Çamyayla, Limestone cliff crevices, 1600 m, 25.viii.1993, Y. Gemici, & Ö. Seçmen 8491 (Holo, EGE).
18. SIDERITIS L. (7: 178)  
e S. akmanii Z.Aytaç, M.Ekici & A. Dönmez, in Flora Mediterranea, 5: 221-224 (1995). [72]  
Type: (Turkey) B3 Afyon: Sandıklı-Şuhut yolu, Kumalar Yaylası, 1520-1550 m, step, 4.viii.1993, Aytaç (6515), Ekici & Dönmez (Holo-GAZI, iso-ANK, HUB).  
e S. caesarea Duman, Aytaç et Başer, in Doğa Tu Botanik D., 22(1): 51-53 (1998). (Sect. Empedoclia (Rafin.) Bentham) [73]  
Type: Turkey. B6 Kayseri: Sakız, Yalak (Yeşilkent), Binboğa Dağı, Tekkekayaşı mevkii, 1800-2200 m, yüksek dağ stebi, 4.viii.1991, Z. Aytaç 4310 - H. Duman (holo-GAZI; iso-ESSE, ANK).
- e S. gulendamii H. Duman & F.A. Karavelioğulları, in Flora Mediterranea, 5: 224-226 (1995). [72]  
Type: (Turkey) B3 Eskişehir: Sivrihisar-Afyon, Aşağıkepen Köyü güneydoğu, jipsli step, 900-950 m, 20.vii.1993, Duman 5219 & Aytaç (Holo-GAZI, iso-ANK, HUB).
- S. scardica Griseb., in Spicil Flor. Rumel. 2: 144 (1844). subsp. scardica [74]  
European Turkey. A1(E) Kırklareli: Demirköy, around Sarpdere village, 800 m, 15.vii.1992, ESSE 9967.
- e S. ozturkii Z. Aytaç et A. Aksoy [75]  
Type: B3 Konya: Derebucak, Kızıldağ, 3 km N of Çamlık town, 1450-1700 m, serpentine rocky place and clearings of *Pinus nigra* forest, 14.vii.1997, Aytaç 7719 & A.Aksoy, para. Aksoy 2312 (holo.Gazi, iso. ANK, HUB).
- e S. vuralii Duman et Başer, in Doğa Tu Botanik D., 22(1): 53-55 (1998) (Sect. Empedoclia (Rafin.) Bentham). [73]  
Type: Turkey. C4 İçel: Anamur, Kazancı-Abanoz, 1300 m, clearing forest and calcareous rocky place, 19.viii.1994, M. Vural 7165, N.Adığüzel & A. Dönmez (holo-GAZI; iso-ESSE, ANK).
19. STACHYS L. (7: 199)  
e S. cydni Kotschy ex Gemici & Leblebici (Sect. Ambleia Bentham), in Doğa Tu Botanik D., 22(5): 359-360 (1998). [76]  
Type: C5 İçel: Arslanköy, Çocakdere, Şahinkaya region, 1560 m, crevices of calcareous rocks, 7.vi. 1995, Gemici 16079 (Holo and Iso EGE); İçel: Çamlıyayla, Payam, Manastır area rocky calcareous crevices, 1450 m, 26.vii.1997, Gem. 8730 (seed).
21. NEPETA L. (7: 264)  
e N. anamurensis Gemici & Leblebici, in Candollea, 50(1): 50 (1995). [14]  
Type: (Turkey) C4 Antalya: Anamur, Abanoz Yaylası, limestone cliff crevices, 1400 m, 25.viii.1993, Y. Gemici & Ö. Seçmen 8498 (Holo, EGE).
- N. humilis Benth., in DC., Prodr., 12: 382 (1858). [77]  
E. Anatolia. C10 Hakkari: Yüksekova-Varağöz, 2150 m, 1.vii.1983, wet places, Koyuncu 6263 & Coşkun (AEF and GAZI).

## 28. ORIGANUM L. (7: 297)

e *O. x adanense* Bašer et Duman, in Doğa Tu Botanik D., 22(1): 55 (1998). (*O. bargyli* Mouterde x *O. laevigatum* Boiss.) [73]

Type: Turkey. C6 Osmaniye: Amanos Dağı, Yarpuz-Ağulu, 1190 m, *P. nigra* forest, 20.vii.1995, H.Duman 5881, K.H.C. Bašer and A. Altıntaş (holo-GAZI; iso-ESSE, ANK).

e *O. husnucan-baserii* H. Duman, Z. Aytaç & A. Duran, in Flora Mediterranea, 5: 226-228 (1995). [72]

Type: (Turkey) C4 Antalya: Alanya, Çökele-Gökbel, *Pinus nigra* açığı, kalkerli alanlar, 1350 m, 9.viii.1994, Duman 5602 Aytaç & Duran (Holo-GAZI, iso-ANK, HUB, ESSE, GAZI).

## 36. THYMUS L. (7: 349)

*T. syriacus* Boiss., in Boiss. Diagn. Ser., 1, 12. p.47. [74]

S.E. Anatolia. C6 Gaziantep: Beylerbeyi village around *Pistachia* plantations, 10.viii.1993, ESSE 10087.

## 39. MENTHA L. (7: 384)

*M. x dumetorum* Schultes, in Obs. Bot. 108 (1809). (*M. aquatica* L. x *M. longifolia* (L.) Hudson.) [78]

N.W. Anatolia. A3 Sakarya: Between Sakarya and Karasu, around Limandere, waterside, 70 m, 29.ix.1995, G. Tarımcılar, BULU 9428. A3 Bolu: In vicinities of the Abant Lake, 1350 m, 1.x.1995, G. Tarımcılar, BULU 9431. A4 Zonguldak: Between Karabük and Eskipazar, around Çay village, streamside, 1450 m, 21.viii.1994, G. Kaynak and G.Tarımcılar, BULU 8837; Safranbolu, 1.5 km from Safranbolu to Akören village, damp and shady places, 1310 m, 22.viii.1994, G. Tarımcılar and G. Kaynak, BULU 8899; Ereğli, parts of Kızılçapınar, streamside, 40 m, 30.ix.1995, G. Tarımcılar, BULU 9429; Hisarönü, around Şahnalar village, 20 m, 30.ix.1995, G. Tarımcılar, BULU 9430; Türkeli, around Ömerdüzü village, 310 m, 26.viii.1994, G. Kaynak and G. Tarımcılar, BULU 9048. A4 Kastamonu: Çiftlik, 1090 m, roadside, 23.viii.1994, G. Tarımcılar and G. Kaynak, BULU 8955. A8 Rize: Çamlıhemşin, roadside, 440-520 m, 28.viii.1993, G. Kaynak and G. Tarımcılar, BULU 7875-7876.

## 42. SALVIA L. (7: 400)

e *S. aytachii* M.Vural & N. Adığuzel, in Doğa Tu Botanik D., 20(6): 531-533 (1996). [79]

Type: (Turkey) B3 Ankara: Polatlı-Sivrihisar, 18. km Açıkır, 840 m, 6.vi.1990, Aytaç 3071, Duman & Adığuzel (holo.: GAZI; iso.: ANK).

## 103. ARISTOLOCHIACEAE (7: 551)

## 2. ARISTOLOCHIA L. (7: 552)

e *A. baseri* Malyer et Erken, in Doğa Tu Botanik D., 21(6): 381-382 (1997). [80]

Type: (Turkey) C2 Antalya: Elmalı: Sütleğen, Kazanpınar mevkii, 1300 m, 21.v.1993, H.Malyer ESSE 11592 (Holo ESSE, ISO, BULU).

## 104. EUPHORBIACEAE (7: 566)

## 5. EUPHORBIA L. (7: 571)

*E. nutans* Lag., in Gen. et Sp. Nov. 17 (1816). [81]

S.E. Anatolia. C5 Adana: Ceyhan-Sirkeli village in corn field, 3.viii.1995 (O.G. 1062) 40 m, O. Gönen. C5 Adana: Kozan and C6 Adana: Kadırli districts in cotton, corn and soybean fields, 1996, 30-40 m, O. G. obs.

## \* FLUEGGEA Willd.

e *F. anatolica* Gemici, in Edinb. Jour. Botany, 50(1): 75 (1993). [82]

Type: (Turkey) C5 Mersin (İçel): N of Tarsus, valley of Kadıncık II Dam, W of the dam, limestone slopes and among *Ceratonia siliqua* trees, 340 m, 14.vi.1990, Y.Gemici 6330 (holo. EGE, iso.E).

## 115. BETULACEAE (7:688)

## 2. ALNUS Miller (7: 691)

*A. glutinosa* (L.) Gaertn., in Fruct. & Sem., 2: 54, t. 90 (1790).

e subsp. *betuloides* Anşin, in Karaca Arboretum Magazine, 2: 48-49 (1993). [83]

Type: (Turkey) B8 Erzurum: Hınıs, environment of Acar (Şıhbirdere) and Hisarcık villages, 1500-1750 m, by streams, 10.vi.1990, Z. Özder (KATO.9886).

## 116. SALICACEAE (7: 694)

## 1. SALIX L. (7: 694)

*S. myrsinifolia* Salisb., in Stirp. Allert.: 394 (1796). [84]

N. Turkey. A4 Kastamonu: N of İlgaç mountains, Dibeyrek valley, *Abies bornmuelleriana* forests, river banks, 1750 m, 18.viii.1992, M. Avcı, s.n., *ibid.*, 15.viii.1995, M. Avcı, ISTE 27492! *ibid.*, 15.vi.1997, M.Avcı, ISTE 27607! ISTE 74659! KOR 39201, 39205!

*S. purpurea* L., in Sp. Pl. 1017 (1753).

e subsp. *leucodermis* Yalt., in Jour. of Facul. of Forestry of Ist. Univ., 38(2): 95 (1988). [85]

Holotype: (Türkiye) C2 Denizli: Kale, Eskere, Çiçekli Büyük Kızıldağ (Sandras Dağı), Yumaklı Uluca Deresi,

1200-1400 m, *Pinus nigra* ormanı, 23.v.1989, F. Yaltırık (ISTO 26660); massif of Sandras Dağı (Çiçekli Büyük Kızıldağ), stony bank of Uluca Dere, 1400-1500 m, 29.vi.1989, Borantynski, Karaca, Zielinski, KOR 516, 517.

118. RUBIACEAE (7:722)

6. GALIUM L. (7:767)

e *G. aladaghense* Parolly, in Willdenowia, 25: 244-246 (1995). [60]

Type: (Turkey) C5 Niğde, Aladağları, Yalak Deresi, 37°50' N, 35°10' E, 2640 m, steinschuttflur mit *Lamium eriocephalum* und *Anthriscus kotschyi*, 45°ENE-exp, 24.viii.1992, Parolly A 836 (B).

*G. recurvum* Req. ex DC., in Prodr., 4: 609 (1830). [86]

W. Anatolia. B1 Balıkesir: Ayvalık, NNE of Alibey Island, Çataltepe, igneous rock crevices, 40 m, 9.v.1995, K. Alpınar, ISTE 67870; W of Alibey Island, Karaada, southern point, 20 m, 30.iv.1997, K. Alpınar, ISTE 74047.

MONOCOTYLEDONES

135. LILIACEAE (8: 67)

9. ASPHODELINE Reichb. (8:88)

e *A. sertachae* Tuzlacı, in Candollea 53: 424 (1998). [87]

Type: (Turkey) C4 Antalya: Gazipaşa, Toros Mountains near Çığlık Village, 2 km above Aşarbaşı, 1820 m, 26.vi.1996, E. Tuzlacı MARE 4961 (Holo-, MARE).

e *A. turcica* Tuzlacı, in Candollea 53: 426 (1998). [87]

Type: (Turkey) C3 Antalya: Gebiz, west of Bozburun Mountain, Boğaz Ağızı, Sanlı Beli, 1470 m, 28.vi.1996, E.Tuzlacı MARE 5000 (Holo-, MARE).

11. ALLIUM L. (8: 98)

e *A. anatomicum* N. Özhata & B. Mathew, in Kew Bulletin, 50(4): 723-724 (1995). [88]

Type: (Turkey) C3 Antalya: province, Çobanisa to Korkuteli, 1050 m, 1.vi.1988, N. & E. Özhata, ISTE 58856 (holotype ISTE !).

e *A. enginii* N. Özhata & B. Mathew, in Kew Bulletin, 50(4): 724-726 (1995). [88]

Type: (Turkey) C5 İçel: Mut to Kirobasi, 15.vi.1990, N. & E. Özhata, ISTE 61830 (holotype ISTE !, isotype K !).

e *A. fethiyense* N. Özhata & B. Mathew, in Kew Bulletin, 50(4): 726-728 (1995). [88]

Type: (Turkey) C2 Muğla: province, 28.v.1988, N. & E. Özhata, ISTE 58655 (holotype ISTE !, isotype K !).

*A. hymenorrhizum* Ledep., in Fl. Altaica, 2: 12 (1830). [89]

E. Anatolia. A9 Kars: Göle, 55078, N. Demirkuş.

e *A. karacae* M. Koyuncu (Sect. Scorodon C. Koch), in Karaca Arboretum Magazine, 2(4): 177-179 (1994). [90]

Type: (Turkey) C4 Konya: between Hadim-Taşkent, open forest and step, 1450 m, 22.vii.1993, H. Karaca, T. Ekim, M. Koyuncu and A. Güner (holo. AEF 18200).

e *A. pseudoalbidum* Friesen & Özhata, in Feddes Repertorium 109(1-2): 25-27 (1998). [91]

Type: A9 Kars: Tractus Susuz Hasköy, in itinere ad montem versus fontis (ISTE 33484 !, isotypus GAT !).

e *A. tuncelianum* (Kollmann) N. Özhata, B. Mathew & Şiraneci, in Kew Bulletin, 50(4): 723 (1995). [88] stat. nov.

Type: (Turkey) B7 Tunceli: province, Munzur Dağ, Aksu Dere above Ovacık, 1800 m, 21.vii.1957, Davis 31498 (holotype E, isotype K !).

e *A. turicum* N. Özhata & Cowley (Sect. Codonoprasum), in Kew Bulletin, 49(3): 481 (1994). [92]

Type: (Turkey) B8 Batman (formerly Siirt), Sason, Halkış Dağ, 1152 m, v/vi.1990, steep stony slopes under mixed trees, N. Stevens, 74 A (holo-type K; isotype ISTE).

18. MUSCARI Miller (8: 245)

e *M. anatomicum* Cowley & N. Özhata, in Kew Bulletin, 49(3): 485-487 (1994). [92]

Type: S Turkey. C5 İçel: 17 km from Arslanköy, above Dümbelek Pass, 2150-2200 m, 14.vi.1990, calcareous, open rocky slopes, beside cemetery, Iter Anatolicum 63/90-267 (ISTE 61781, PI 689/90) (holotype ISTE; isotype K).

20. BELLEVALIA Lapeyr. (8: 264)

e *B. anatolica* B. Mathew & N. Özhata, in Kew Bulletin, 49(3): 483-485 (1994). [92]

Type: (Turkey) B7 Elazığ: N. of Elazığ near Pertek, on road to Tunceli, rocky hillside, 4.vi.1987, fl. cult. Kew, 10 April 1990, B. Mathew 11033 (holotype K).

22. LILIUM L. (8: 279)

e L. akkusianum R. Gämperle, in Bulletin of the Alpine Garden Society, 66, 380-382 (1998). [93]

Holotype: A6 Ordu: 25.6.1996, in Anatolia, secundum viam ab Ünye ad Niksar, 6.7 km ab coemeterio Akkuş vici, Niksar versus, in clivo viae, inter frutices fagorum, altitudo 1100-1500 m, leg. René Gämperle. Conservatur in herbario horti botanici turicensis (Z).

### 23. FRITILLARIA L. (8: 284)

e F. sororum J. Persson & K. Persson, in New Plantsman, 5(2): 114-115 (1998). [94]

Type: (Turkey) C5 İçel: Taurus mountains, 500 m, meadow fragments adjoining brook ravine in *Pinus brutia* forest, on limestone, coll. 30.iv.1987, cult. 20.v.1995, J.Persson & K.Persson 87-89 (holo-type, GB).

F. sibthoriana (Smith) Baker, in J.Linn. Soc. (Bot.), 14: 275 (1874).

e subsp. enginiana Byfield & N. Özhatay, in Karaca Arboretum Magazine, 3: 9-11 (1995). [95]

Type: SW Turkey. C2 Muğla: Köyceğiz, Sandras Dağı, Pankuduz Tepe, under woodland, rocky and moist slopes, 1000 m, 6.iv.1977, E.Özhatay 1077 (Holotype ISTE 36685, isotypes ISTO 19211 and MARA 0181).

### 30. COLCHICUM L. (8: 329)

e C. davisii C.D. Brickell, in The New Plantsman, 5(1): 15-18 (1998). [96]

Type: (Turkey) C5 Adana: d. Bahçe Dumanlı Dağ, 1300 m, edge of beech forest, 19.iv.1957, Davis 26938. Cult. R.B.G. Edinburgh, 29.ix.1984 (fl) (Holotypus E; isotypus Wisley).

### 136. AMARYLLIDACEAE (8: 358)

#### 5. GALANTHUS L. (8: 365)

G. byzantinus Baker, in Gard. Chron. Ser.3, 13: 226 (1893).

e subsp. braunerii N. Zeybek, in Doğa Tu Botanik D., 12(1): 101 (1988). [97]

Type: (Turkey) A3 Bolu: Abant Gölü Orman Bölge Şefliği Tomruk Deposu çevresi, 1400 m, N. Zeybek (holo. IZEF 942, iso. ISTE) !

G. byzantinus Baker, in Gard. Chron. Ser.3, 13: 226 (1893).

e subsp. sauerii N. Zeybek, in Doğa Tu Botanik D., 12(1): 101 (1988). [97]

Type: (Turkey) A3 Bolu: Abant Gölü yolu, 1250 m, 12.iv.1978, E. Sauer (holo. BRD. Saarland Univ. Herb. 18643, iso. IZEF) !

G. byzantinus Baker, in Gard. Chron. Ser.3, 13: 226 (1893).

e subsp. tughrului N. Zeybek, in Doğa Tu Botanik D., 12(1): 100 (1988). [97]

Type: (Turkey) A3 İzmit: Keltepe, 1050 m, 8.ii.1985, L.Tuğrul (holo. IZEF 926, iso. ISTE) !

G. elwesii Hook., in Bot. Mag., 101: t. 6166 (1875).

e subsp. akmanii N. Zeybek, in Doğa Tu Botanik D., 12(1): 99 (1988). [97]

Type: (Turkey) A4 Ankara: Uruş, Örencik Köyü, kayalık step yamaçlar, 1500 m, 7.iii.1981, N. Zeybek, M. Koyuncu, M. Aydoğdu (holo. IZEF 409, iso. ISTE) !

G. elwesii Hooker fil., in Bot. Mag., 101: t. 6166 (1875).

e subsp. baytoppii N. Zeybek & E.Sauer, in Doğa Tu Botanik D., 12(1): 96-97 (1988). Syn.: *G.gracilis* Celak. subsp. *baytoppii* N.Zeybek [97] com. nov.

Type: (Turkey) C3 Antalya, Akseki, Çimi Köyü Yaylası, Toptaş Mevkii, ca.1400 m, 6.ii.1984, U. Zeybek (IZEF 1810).

G. elwesii Hook., in Bot. Mag., 101: t. 6166 (1875).

e subsp. melihae N. Zeybek, in Doğa Tu Botanik D., 12(1): 98 (1988). [97]

Type: (Turkey) A3 İzmit: Kaynarca civarı, tarla kenarları, 5.ii.1983, N. Zeybek (holo, IZEF 944, iso, ISTE) !

G. elwesii Hooker, in Bot. Mag., 101: t. 6166 (1875).

e subsp. tuebitaki N. Zeybek, in Doğa Tu Botanik D., 12(1): 98 (1988). [97]

Type: (Turkey) A4 Ankara: Kızılcahamam Milli Park, 1350 m, 16.iii.1983, N. Zeybek (holo. IZEF 912, iso. ISTE) !

G. elwesii Hooker, in Bot. Mag., 101: t. 6166 (1875).

e subsp. yayintaschii N. Zeybek, in Doğa Tu Botanik D., 12(1): 98 (1988). [97]

Type: (Turkey) B2 Manisa: Demirci-Simav (Kütahya) yolu, 4-5 km de tarla içleri, 5.iii.1979, A. Yayıntaş (holo. IZEF 401, iso. ISTE) !

G. elwesii Hook., in Bot. Mag., 101: t. 6166 (1875).

e subsp. wagenitzii N. Zeybek, in Doğa Tu Botanik D., 12(1): 98 (1988). [97]

Type: (Turkey) C5 Mersin: Yavca Köyü, 1250 m, elma

bahçeleri, 11.ii.1982, N. Zeybek (holo. IZEF 905, iso. ISTE) !

*G. fosteri* Bakeri, in Garden. Chron. Ser. 3(5): 458 (1889).

e var. *antepensis*\*, in Türkiye Kardelenleri (*Galanthus* L.), I: 72 (1995). [98] nom. nud.

Type: (Turkey) C6 Gaziantep: Gaziantep-Bahçe, Gaziantep'ten 30 km uzakta, Acarobası Köyü makilikleri, Kalker kayalar arasında humuslu topraklarda, ca.1200 m, 16.ii.1994, N. Zeybek (IZEF 2155).

\* This new taxon not published as valid.

*G. gracilis* Celak, in Sitz.-Ber. Böhm. Ges. Wiss. 1891: 195, t. 9 (1891).

e subsp. *baytoppii* N. Zeybek, in Doğa Tu Botanik D., 12(1): 96 (1988). [97]

Type: (Turkey) C3 Antalya: Akseki, Çimi Köyü Yaylası, Toptaş, 1400 m, 6.ii.1984, U. Zeybek (holo. IZEF 922, iso. ISTE) !

e *G. melihae* (N. Zeybek) E. Sauer & N. Zeybek (*G. elwesii* Hooker subsp. *melihae* N. Zeybek, in Doğa Tu Botanik D., 12(1): 98 (1988) [97], in Türkiye Kardelenleri (*Galanthus* L.), I: 54 (1995). [98]

Type: (Turkey) A3 Kocaeli, Kaynarca civarı ve Adapazarı, Karaçalı Köyü civarlarında çok bol, 5.ii.1983, N. Zeybek (IZEF 1784).

*G. nivalis* L., in Sp. Pl. 288 (1753).

e subsp. *humboldtii* N. Zeybek, in Doğa Tu Botanik D., 12(1): 97 (1988). [97]

Type: (Turkey) A1(E) Kırklareli: İğneada, Eski Orman Kampı, 1-2 m, 19.ii.1982, M.A. Önür (holo. IZEF 906, iso. ISTE) !

*G. nivalis* L., in Sp. Pl. 288 (1753).

e subsp. *subplicatus* N. Zeybek & E. Sauer (syn. *G. plicatus* Bieb. subsp. *subplicatus* N. Zeybek in Doğa Tu Botanik D., 12(1): 99 (1988) [97], in Türkiye Kardelenleri (*Galanthus* L.), I: 84 (1995). [98] com. nov.

Type: (Turkey) A2(E) İstanbul: Bahçeköy, Belgrat Ormani, Kurtkemer, *Carpinus betulus* ağaçları altı, 31.iii.1980, N. Zeybek (IZEF 1833).

e *G. peshmenii* A.P. Davis & C.D. Brickell, in The New Plantsman, 1(1): 17 (1994). [99]

Type: S.W. Turkey, Prov. C3 Antalya: Kemer, Kesmeboğazi-Gedelma Köyü arası, c.300 m, maki-orman topluluğu, 3.xi.1978, Peşmen, Yıldız et Güneş 4125 (holotypus HUB, isotypus E).

*G. plicatus* M. Bieb., in Fl. Taur.-Cauc. 2: 255 (1819).

e subsp. *karamanogluensis* N. Zeybek, in Doğa Tu Botanik D., 12(1): 100 (1988). [97]

Type: (Turkey) A4 Ankara: Kızılcahamam, Akyarma Geçidi, 1550 m, 15.iv.1973, K. Karamanoğlu, M. Koyuncu (holo. AE 4708, iso. IZEF) !

*G. plicatus* M. Bieb., in Fl. Taur.-Cauc. 2: 255 (1819). e subsp. *sublicatus* N. Zeybek, in Doğa Tu Botanik D., 12(1): 99 (1988). [97]

Type: (Turkey) A2(E) İstanbul: Bahçeköy, Belgrat ormanı, Kurt Kemer, *Carpinus betulus* ağaçları altı, 31.iii.1980, N. Zeybek (holo. IZEF 406, iso. ISTE) !

*G. plicatus* M. Bieb., in Fl. Taur.-Cauc. 2: 255 (1819). e subsp. *vardarii* N. Zeybek, in Doğa Tu Botanik D., 12(1): 100 (1988). [97]

Type: (Turkey) C8 Trabzon: Sürmene, Arpalı Köy, 24.iii.1983, N. Zeybek (holo. IZEF 917) !

137. IRIDA CEAE (8: 381)

4. CROCUS L. (8: 413)

*C. biflorus* Miller, in Gard. Dict. ed. 8, no 4 (1768).

e subsp. *albo-coronatus* H. Kerndorff, in Herbertia, 49(1-2): 79 (1993). [100]

Type: (Turkey) Central Taurus Mountains, H. Kerndorff HKEP 9009 (holotypus, K).

*C. biflorus* Miller, in Gard. Dict. ed. 8, no 4 (1768).

e subsp. *fibroannulatus* H. Kerndorff & E. Pasche, in Linzer Biol. Beitr. 29(1): 594-595 (1997). [101]

Type: (Turkey) A8 Artvin: Provinz Artvin, H. Kerndorff & E. Pasche, HKEP 9361 (holo. K).

*C. biflorus* Miller, in Gard. Dict. ed. 8, no 4 (1768).

e subsp. *wattiorum* B. Mathew, in The New Plantsman, 2(3): 183-184 (1995). [102]

Type: SW Turkey, C3 Antalya: province, Tahtalı Dağ, 6.xi.1986, P. & P. Watt s.n.; cult. B. Mathew, florendi 20.x.1994 (holotypus K).

e *C. kerndorffiorum* E. Pasche, in Herbertia, 49(1-2): 67-68 (1993). [103]

Type: (Turkey) Central Taurus Mountains, E. Pasche HKEP 9010 (holotypus, K).

e *C. mathewii* Kerndorff & Pasche, in The New Plantsman, 1(2): 102-105 (1994). [104]

Type: (Turkey) Antalya: Lycian Taurus Mountains, 400-1100 m, 16.xi.1992, HKEP 9291 (holotype, K).

e *C. paschei* H. Kerndorff, in *Herbertia*, 49(1-2): 76-77(1993). [100]

Type: (Turkey) Eastern Taurus Mountains, H. Kerndorff HKEP 9034 (holotype, K)

144. CYPERACEAE (9: 32)

1. *CYPERUS* L. (9: 34)

*C. glomeratus* L., in *Cent. Pl.*, 2: 5 (1756). [105]

European Turkey. A1(E) Edirne: Uzunköprü, Çakmak köyü, çeltik tarlası, 28.viii.1995, S. Sönmez, ISTE 68563.

145. GRAMINEAE (Poaceae) (9: 158)

23. *BROMUS* L. (9: 272)

*B. tomentellus* Boiss. (Sect. *Pnigma* Dum.), in *Diagn. Pl. Or. Nov.*, 1(7): 126 (1846).

subsp. *nivalis* (Bornm.) H. Scholz et Byfield, in *Doğa Tu Botanik D.* (1998). [106] stat. nov.

N.E. Anatolia. A9 Çoruh: 23 km from Şavşat to Ardahan, before Çam pass, 2350 m, alpine pasture grassland on igneous rocks, abundant, 4.vii.1994, A.J. Byfield & D. Pearman B 1289 (ISTE 67154).

75. *POA* L. (9: 470)

e *P. akmanii* Soreng, P.Hein & Scholz, in *Willdenowia*, 27: 196-197 (1997). [107]

Holotype: Turkey, prov. C2 Isparta: Isparta, BarlaDağı, Gelincikdağ, ca.5 km W of Barla (W of Eğirdir Gölü and c.35 km NE of Isparta) 38° 08' N, 30° 45' E, 2287 m, alpine, tall, shady, NE facing limestone cliffs, on mossy ledges protected from marauding goats, above snow field, 21.vii.1993, R.J. Soreng, J.I. Davis, K. Güney & Ü. Bingöl 4140 (US; iso: ANK, B, E, ISTE, LIV).

P. *asiae-minoris* H.Scholz & Byfield, in *Doğa Tu Botanik D.* –baskıda-. [106]

Holotype: N.W.Anatolia. A3 Bolu: Abant Lake, S.W. end of lake, 1450 m, wet peat fen, extremely local, 14.vi.1995, A.J. Byfield & R. Fitzgerald B 2351 (ISTE 69705).

116. *SPOROBOLUS* R.Br. (9: 581)

*S. fertilis* (Steud.) W.D. Clayton, in *Kew Bull.* 19(2): 291 (1965). [59]

N.E. Anatolia. A8 Artvin: Kemalpaşa, beach W. of river mouth, sandy dry grassland behind beach, 5 m, 17.viii.1995, A.J.Byfield and S.Atay (ISTE 69298). Det. T.Cope (Kew).

Alien species.

\* *MIBORA* Adanson

*M. minima* (L.) Desv., in *Obs. Pl. Angers* 48 (1818). [41]

European Turkey. A1(E) Edirne: Pehlivanköy, elevated commonland a few kilometres from Pehlivanköy towards Kircasalih, 25.ii.1994, A.J.Byfield et al. (AJB 295), det. A.J.Byfield, conf. T.A.Cope, ISTE 66249.

\* *MICROSTEGIUM*

*M. vimineum* (Trin.) A.Camus, in *Ann. Soc. Linn. Lyon*, 68: 201 (1921). [106]

(Turkey). N.E. Anatolia. A7 Giresun: Espiye River, immediately E. of Espiye town, wet, seasonally-flooded river margin on gravels & sands, 5m, 11.x.1994, A.J. Byfield B 1334 (ISTE 67575).

Additional List

13. CARYOPHYLLACEAE (2:15)

13. SPERGULARIA (Pers.) J. & C.Presl (2:93) [108]

*S. sezer-zenginii* Yıldırımlı, in *Ot Sistematisk Botanik Dergisi*, 4(2): 4 (1997).

Type: Turkey. B9 Ağrı: Hamur, Murat river, sandy places, 1650 m, 2.vii.1998, Ş.Yıldırımlı 23037 (Holo: Herbarium Yıldırımlı, iso: GAZİ, HUB, UPA).

45. LEGUMINOSAE (3:1)

40. *VICIA* L. (3:274)

*V. erzurumica* N.Demirkuş & S. Erik (sect.*Cracca* S.F.Gray), in *Willdenowia* 28: 154 (1998). [109]

Holotypus: (Turkey). A8 Erzurum. Tortum, Azort Yayları, orman yolu, taşlık alanlar, 1900 m, 20.vii.1995, Demirkuş 4481 (HUB: isotypi: GAZI, B).

63. UMBELLIFERAE (4: 265)

10. CHAEROPHYLLUM L.(4:312)

*C. posofianum* S. Erik & N. Demirkuş, in *Willdenowia* 28: 153 (1998). [109]

Holotypus: (Turkey) A9 Ardahan (Kars): Posof, fields between Al village and Posof river, 1500-1700 m, 27.vii.1985, Demirkuş 2960 (HUB).

70. COMPOSITAE (5:1)

79. *CENTAUREA* L. (5:465)

*C. hadimensis* Wagenitz, K.Ertuğrul & H.Dural, in *Willdenowia* 28: 157 (1998). [110]

Holotypus: (Turkey). C4 Konya: Konya-Hadim, Gevne Valley, SW of Tosmur Yurdu, rocky places, 1500 m,

30.vi.1996, Ertuğrul & Dural 1715 (KNYA: isotypus: GOET).

#### CARDUNCELLUS

*C. caeruleus* (L.) C. Presl, in *Flora Sicula*, 1:30 (1826).

var. *incisus* DC., in *Prodr.*, 6: 615 (1837). [111]

W. Anatolia. B1 Balıkesir: Ayvalık, Alibey adası, Patriça civarı, 10 m, 7.v.1996, K. Alpinar, ISTE 71692! Pınar Adası'nın doğu kıyısı, hafif eğilimli, güneşli yerler, 5 m,

10.v.1996, K. Alpinar, ISTE 71768! Dolap Adası-Mitralöz Burnu arası, d.s., 26.iv.1997, K. Alpinar, ISTE 73872! Alibey Adası, Alibey Tepe, *Pinus brutia* orman açıklığı, 120 m, 26.v.1997, K. Alpinar, ISTE 74209!

#### Acknowledgement

We are grateful to the following botanists for their valuable contribution: Doç. Zeki Aytaç, Mr. Andrew Byfield, Doç. Hayri Duman, Prof. Tuna Ekim

#### References

1. Davis, P.H. (ed.), *Flora of Turkey and the East Aegean Islands*, Vol. 1-9, University Press, Edinburgh (1965-1985).
2. Davis, P.H. (ed.), *Flora of Turkey and the East Aegean Islands (Supplement)*, Vol. 10, University Press, Edinburgh (1988).
3. Özhatay, N., Kültür, ş., Aksoy, N., Check-List of additional taxa to the supplement Flora of Turkey, *Doğa Tu Botanik D.*, 18, 497-514 (1994).
4. Browicz, K., An unpublished Flora of Constantinople from the second half of the XIX Centur, in Davis & Hedge, *Festschrift, Plant Taxonomy Phytogeography and related subjects* (ed. Kit Tan), Edinburgh University Press, Edinburgh (1989).
5. Alpinar, K., Some contributions to the Turkish Flora, *Edinb. Jour. Bot.*, 51, 1, 65-73 (1994).
6. Boissier, P.E., *Flora Orientalis*, 5, 749, Basileae, Geneva & Lugduni (1884).
7. Altınayar, G., Su Yabancıotları, T.C.Bayındırılık ve İskan Bakanlığı, Devlet Su İşleri Genel Müdürlüğü, İşletme ve Bakım Dairesi Başkanlığı, Ankara (1988).
8. Ünal, M., Üzen, E., A new aquatic fern record for the flora of Turkey: *Azolla filiculoides* Lam., *Doğa Tu Botanik D.*, 20, 4, 379-381 (1996).
9. Yaltırık, F., Boydak, M., Ülkemizde yeni bir kızılçam varyetesi ve Kızılçamın varyasyonları, I.Ü. Orman Fak. Derg., Seri A, Cilt 39, 2, 42-64 (1989).
10. Boydak, M., Türkiye'de Anadolu Karaçamının yeni bir varyetesi, I. Ü. Orman Fak. Derg. Seri A, Cilt 39, 1, 119-129 (1989).
11. Baytop, A., Quelques notes sur la flore de la Turquie d'Europe, IV (Trakya Florası üzerinde bazı notlar, IV), Istanbul Ecz. Fak. Mec., 8, 109-114 (1972).
12. Özhatay, N., Özhatay, E., A new white *Paeonia* L. from NW Turkey: *P. mascula* Miller subsp.*bodurii* N. Özhatay, *The Karaca Arboretum Magazine*, 3, 1, 17-26 (1995).
13. Liden, M., New taxa of tuberosus *Corydalis* (Fumariaceae), *Willdenowia*, 26, 23-35, Berlin (1996).
14. Gemici, Y., Leblebici, E., Seven new species for the Flora of Turkey, *Candollea*, 50, 1, 41-50 (1995).
15. Duman, H., A new species from South-Anatolia - *Aethionema alanyae* (Brassicaceae), *The Karaca Arboretum Magazine*, 2, 4, 171-176 (1994).
16. Vural, M., Adıgüzell, N., A new species from Central Anatolia-*Aethionema dumanii* (Brassicaceae), *Doğa Tu Botanik D.*, 19, 4, 481-483 (1995).
17. Ertuğrul, K., Beyazoğlu, O., A new species from South Anatolia - *Aethionema karamanicus* (Cruciferae), *Doğa Tu Botanik D.*, 21, 2, 99-101 (1997).
18. Orcan, N., Dudley, T.R., Misirdal, H., Two new taxa of *Alyssum* for the Flora of Turkey, *Bot. Chron.*, 11, 59-62 (1994).
19. Carlström, A., A survey of the Flora and phytogeography of Rodhos, Simi, Tilos and the Marmaris Peninsula (SE Greece, SW Turkey), Lund 1987, (ISBN 91.7970.096.1).
20. Yıldırımlı, ş., *Lunaria* (Brassicaceae) a new genus record for the flora of Turkey, *Ot Sistematisk Botanik Dergisi*, 3, 1, 65-68 (1996).
21. Yıldırımlı, ş., *Viola bocquetiana* (Violaceae) a new species from Turkey, *Ot Sistematisk Botanik Dergisi*, 1, 1, 1-4 (1994).
22. Duman, H., A new species of *Minuartia* L. (Caryophyllaceae) from south Anatolia, *The Karaca Arboretum Magazine*, 4, 4, 159-162 (1998).
23. Burton, R.M., A new variety of *Dianthus elegans* D'Urv., *The Karaca Arboretum Magazine*, 3, 4, 183-186 (1997).
24. Özhatay, N., Alpinar, K., A new record for the Turkish Flora: *Gypsophila glomerata* Pallas ex Bieb., *Istanbul Ecz. Fak. Mec.*, 16, 10-13 (1980).
25. Duman, H., Aytaç, Z., Vural, M., Two new taxa from South Anatolia, *Doğa Tu Botanik D.*, 19, 4, 477-479 (1995).
26. Aytaç, Z., A new species of *Silene* (Caryophyllaceae) from Anatolia (Turkey), *Thaiszia*, J.Bot., Kosice, 8, 7-11 (1998).

27. Leblebici, E., Duman, H. and Aytaç, Z.. *Polygonum ekimianum* (*Polygonaceae*), a new species from Anatolia, *Willdenowia*, 23, 163-166 (1993).
28. Yıldız, B., *Rumex arifolius* All. (*Polygonaceae*), new record for the Flora of Turkey, *Doğa Tu Botanik D.*, 19, 5, 567-568 (1995).
29. Yıldırımli, Ş.. *Ceratocarpus turkestanicus* Sav.-Rycz. (*Chenopodiaceae*), a new record for the flora of Turkey, Dobrudscha (Bulgaria-Romania), and Crimea (Ukraine), *Ot Sistematisk Botanik Dergisi*, 3, 2, 99-104 (1996).
30. Freitag, H., Özhatay, E.. A new subspecies of *Salsola canescens* (*Chenopodiaceae*) from SW Anatolia, Turkey, *Willdenowia* 27, 185-190 (1997).
31. Browicz, K., Chorology of Trees and Shrubs in South-West Asia and Adjacent Regions, Polish Academy of Sciences, Institute of Dendrology, Kornik, 10, 34 (1994).
32. Yıldırımli, Ş.. *Hypericum peshmenii* (*Hypericaceae*) a new species from Turkey, *Ot Sistematisk Botanik Dergisi* 2, 2, 11-14 (1995).
33. Güner, A., Vural, M., Duman, H., Dönmez, A.A., Şağban, H., The Flora of the Köyceğiz-Dalyan specially protected Area (Muğla-Turkey), *Doğa Tu Botanik D.*, 20, 329-371 (1996).
34. İlarslan, R., Yurdakulol, E., A new species of *Erodium* (*Geraniaceae*) from Turkey, *Flora Mediterranea* 4, 167-170 (1994).
35. Duman, H., Başer, K.H.C., Malyer, H., A new species of *Chamaecytisus* Link (*Leguminosae*) from SW Anatolia, *The Karaca Arboretum Magazine*, 4, 1, 13-18 (1997).
36. Malyer, H., A new record for the Flora of Turkey, *Doğa Tu Botanik D.*, 20, 5, 473-475 (1996).
37. Podlech, D., Aytaç, Z.. Two new species of *Astragalus* L. sect. *Dissitiflori* DC. (*Fabaceae*) from Turkey, *Sendtnera*, 5, 265-268 (1998).
38. Duman, H., Ekim, T., *Astragalus serpentinicola* (A. sect. *Sisyphorus*, *Leguminosae*), a new species from SW Turkey, *Willdenowia*, 27, 181-184 (1997).
39. Ladizinsky, G., A new species of *Lens* from south-east Turkey, *Bot. Journal of Linnean Society*, 123, 257-260 (1997).
40. Ertekin, A.S.. A new record for the Flora of Turkey, *Doğa Tu Botanik D.*, 18, 1, 33-34 (1994).
41. Byfield, A., Özhatay, N.. Two new species to Turkey from Turkish Thrace: *Mibora minima* (L.) Desv. (*Gramineae*) & *Trifolium ornithopodoides* (L.) Sm. (*Leguminosae*), *Doğa Tu Botanik D.*, 22, 425-429 (1998).
42. Kalheber, H., The Genus *Alchemilla* L. (*Rosaceae*) in the Turkish Vilayet Rize (Northeastern Anatolia) with some remarks on the distribution of the genus in other parts of Northern Anatolia, *Sendtnera*, 2, 389-430 (1994).
43. Hayırlioğlu, S., Beyazoğlu, O.. Five new *Alchemilla* (*Rosaceae*) records for the Flora of Turkey, *Doğa Tu Botanik D.*, 21, 309-312 (1997).
44. Hayırlioğlu-Ayaz, S., Beyazoğlu, O.. A new species of *Alchemilla* (*Rosaceae*) from Turkey, *Ann. Bot. Fennici*, 34, 109-113 (1997).
45. Güner, A., Duman, H., Critics on *Pyrus boissieriana* Buhse subsp.*crenulata* Browicz, *The Karaca Arboretum Magazine*, 2, 4, 165-170 (1994).
46. Leblebici, E., Seçmen, Ö.. A new species of *Lythrum* (*Lythraceae*) from the Western Black Sea Region of Turkey, *Doğa Tu Botanik D.*, 19, 5, 559-560 (1995).
47. Göktürk, R.S., Sümbül, H., A new record for the flora of Turkey: *Ludwigia peploides* (Kunth) P.H.Raven, *The Karaca Arboretum Magazine*, 4, 3, 109-112 (1998).
48. Duman, H., Güner, A., A new record for the flora of Turkey, *Doğa Tu Botanik D.*, 20, 4, 383-384 (1996).
49. Metzger, J.. *Sedum aytacianum* (*Crassulaceae*), a new species from Turkey, *Cactus and Succulent Journal (U.S.)*, 66, 259-261 (1994).
50. 't Hart, H., Variation in *Sedum litoreum* and the *S. rubens* complex (*Crassulaceae*); new names and combinations, *Ot Sistematisk Botanik Dergisi*, 2, 2, 1-9 (1995).
51. Metzger, J.. *Sedum vuralianum* (*Crassulaceae*), a new species from N.E.Anatolia, *British Cactus & Succulent Society Journal*, 11, 4, 100-102 (1993).
52. Özhatay, N., Özhatay, E.. A new record for the Turkish Flora: *Saxifraga graeca* Boiss. et Heldr., *Istanbul Ecz. Fak. Mec.*, 16, 14-16 (1980).
53. Yıldız, B., Bahçecioğlu, Z.. A new species from Malatya/Turkey: *Echinophora lamondiana* B.Yıldız & Z. Bahçecioğlu (*Apiaceae/Umbelliferae*), *The Karaca Arboretum Magazine*, 4, 1, 7-12 (1997).
54. Tanker, N., Koyuncu, M., Coşkun, M., Güvenç, A., Özgen, U., Idris Dağı (Ankara) Bitkileri, *Ankara Ecz. Fak. Der.*, 22, 1-2, 1-20 (1993).
55. Pimenov, M. & Kluykov, V.E.. A new monotypic section in *Prangos* Lindl. (*Umbelliferae*) and the correct name for its species, *Candollea*, 52, 1, 109-111 (1997).
56. Dawud Al-Eisawi, F.L.S. & Stephen L., Jury, F.L.S., A taxonomic revision of the genus *Tordylium* L. (*Apiaceae*), *Bot. Journal of Linnean Society*, 97, 357-403 (1988).
57. Karamanoğlu, K., Koyuncu, M., *Türkiye Valeriana* (Kediotu) türleri üzerinde sistematik araştırmalar, *Ankara Ecz. Fak. Der.*, 4, 1, 149-178 (1974).

58. Göktürk, S., Sümbül, H., A new species of *Cephalaria* (*Dipsacaceae*) from South Anatolia, Turkey, Ann. Bot. Fennici, 34, 153-155 (1997).
59. Byfield, A., Baytop, A., Three alien species new to the flora of Turkey, Doğa Tu Botanik D., 22, 3, 205-208 (1998).
60. Parolly, G., New taxa and noteworthy records from the Western and Middle Taurus Range, Turkey, Willdenowia, 25, 239-252 (1995).
61. Baytop, A., Byfield, A.J., The Presence of *Logfia minima* (Sm.) Dumort. (*Compositae*) in Turkey, Doğa Tu Botanik D., 21, 4, 245-246 (1997).
62. Tuzlaci, E., A new record for the flora of Turkey: *Onopordum sirsangense* Rech. f., Mar. Üniv. Ecz. Fak. Derg., 2, 1, 57-61 (1986).
63. Burton, R.M., A new thistle species from south west Turkey, The Karaca Arboretum Magazine, 3, 3, 123-126 (1996).
64. Gottschlich, G., *Hieracium nydeggerorum* Gottschlich, eine neue *Hieracium*-Art aus der Türkei, Bauhinia, 11, 4, 239-245 (1996).
65. Davis, P.H., Sorger, F., A new *Campanula* from Anatolia, Notes RBG, 37, 2, 265-266 (1979).
66. Güner, A., A new record for the Flora of Turkey and a new subspecies from Anatolia, Candollea, 39, 1, 345-348 (1984).
67. Dönmez, A.A., Güner, A., A new species from Central Anatolia: *Campanula kirikkaleensis* (*Campanulaceae*), The Karaca Arboretum Magazine, 2, 3, 115-119 (1993).
68. Anşin, R., Terzioğlu, S., Mor çiçekli orman gülünün yeni bir varyetesi (*Rhododendron ponticum* L. subsp.*ponticum* var.*heterophyllum* Anşin, var.*nova*), Doğa Tr. J. of Agricultural and Forestry, 18, 137-140 (1994).
69. Yıldırımlı, Ş., *Buglossoides glandulosum* (*Boraginaceae*) a new record for the Flora of Turkey, Ot Sistematič Botanik Dergisi, 1, 1, 57-60 (1994).
70. Çakan, H., Düzenli, A., A new record for the Flora of Turkey, Doğa Tu Botanik D., 19, 4, 491-492 (1995).
71. Duman, H., A new species of *Teucrium* L. (*Labiatae*) from SW Anatolia, The Karaca Arboretum Magazine, 4, 3, 125-130 (1998).
72. Duman, H., Aytaç, Z., Ekici, M., Karavelioğulları, F.A., Dönmez, A., Duran, A., Tree new species (*Labiatae*) from Turkey, Flora Mediterranea, 5, 221-228 (1995).
73. Duman, H., Başer, K.H.C., Aytaç, Z., Two new species and new hybrid from Anatolia, Doğa Tu Botanik D., 22, 1, 51-57 (1998).
74. Başer, K.H.C., Vural, M., et al., Two new records for the Flora of Turkey, Doğa Tu Botanik D., 19, 4, 489-490 (1995).
75. Aytaç, Z., Aksoy, A., A new *Sideritis* L. species (*Labiatae*) from Turkey, Flora Mediterranea (1998) –baskıda-
76. Gemici, Y., Leblebici, E., A new species from Southern Anatolia: *Stachys cydni* Kotschy ex Gemici & Leblebici, Doğa Tu Botanik D., 22, 5, 359-362 (1998).
77. Aytaç, Z., Yıldız, G., A new record for the Flora of Turkey, Doğa Tu Botanik D., 20, 4, 385-386 (1996).
78. Tarımcılar, G., Kaynak, G., A new record for the Flora of Turkey, Doğa Tu Botanik D., 21, 4, 247-249 (1997).
79. Vural, M., Adıgüzel, N., A new species from Central Anatolia: *Salvia aytachii* M. Vural & N. Adıgüzel (*Labiatae*), Doğa Tu Botanik D., 20, 6, 531-534 (1996).
80. Malyer, H., Erken, S., A new species from Turkey *Aristolochia baseri* (*Aristolochiaceae*), Doğa Tu Botanik D., 21, 6, 381-383 (1997).
81. Gönen, O., Uygur, F.N., A new record for the Flora of Turkey: *Euphorbia nutans* Lag., Türkiye Herboloji Dergisi, 1, 2, 14-17 (1998).
82. Gemici, Y., A new species of *Flueggea* (*Euphorbiaceae*) from Anatolia, Edinb. Jour. Botan., 50, 1, 75-77 (1993).
83. Anşin, R., Özder, Z., A new taxon of black alder *Alnus glutinosa* subsp.*betuloides* (*Betulaceae*), The Karaca Arboretum Magazine, 2, 2, 47-51 (1993).
84. Avci, M., Zielinski, J., *Salix myrsinifolia* Salisb. (*Salicaceae*) a new species for the flora of Turkey, The Karaca Arboretum Magazine, 4, 2, 49-54 (1998).
85. Yaltırık, F., Türkiye Florası için yeni bir söğüt taksonu: *Salix purpurea* L. subsp.*leucodermis* Yalt. subsp. nova, I.Ü. Orman Fak. Der. Seri A, 38, 2, 92-95 (1988).
86. Alpinar, K., A new taxon for Turkey: *Galium recurvum* Req. ex DC. (*Rubiaceae*), Doğa Tu Botanik D., 22, 4, 295-298 (1998).
87. Tuzlaci, E., Revision of the genus *Asphodeline* (*Liliaceae*). II. Two new species from Turkey, Candollea, 53, 423-433 (1998).
88. Özhatay, N., Mathew, B., New taxa and notes on the genus *Allium* (*Alliaceae*) in Turkey and Arabia, Kew Bulletin, 50, 4, 723-731 (1995).
89. Özhatay, N., Allium in Turkey: distribution, diversity, endemism and chromosome number. Proceedings of the 5th Optima Meeting, İstanbul, 8-15 September 1986 (eds.. N.Özhatay, H.Demiriz), İstanbul, 247-271 (1993).
90. Koyuncu, M., A new species of *Allium* L. (*Liliaceae*) from Southern Anatolia, The Karaca Arboretum Magazine, 2, 4, 177-180 (1994).
91. Friesen, N., Özhatay, N., New taxa and notes on *Allium* subgenus *Rhizirideum* (*Alliaceae*) in Turkey, Feddes Repertorium 109, 1-2, 25-31 (1998).

92. Cowley, J., Özhatay, N., Mathew, B., New species of *Alliaceae* et *Hyacinthaceae* from Turkey. *Kew Bulletin*, 49, 3, 481-489 (1994).
93. Gämperle, R., A new species of lily from Turkey. *Bulletin of the Alpine Garden Society*, 66, 378-389 (1998).
94. Persson, J., Persson, K., *Fritillaria sororum* (*Liliaceae*): a new species from Turkey. *The New Plantsman*, 5, 2, 114-116 (1998).
95. Özhatay, N., Byfield, A., A new *Fritillaria* L. (*Liliaceae*) from southwest Turkey. *The Karaca Arboretum Magazine*, 3, 1, 7-16 (1995).
96. Brickell, C., *Colchicum davisii* (*Colchicaceae*) a new species from Turkey. *The New Plantsman*, 5, 1, 15-22 (1998).
97. Zeybek, N., Taxonomical investigations on Turkish Snowdrops (*Galanthus* L.). *Doğa Tu Botanik D.*, 12, 1, 89-103 (1988).
98. Zeybek, N., Sauer, E., *Türkiye Kardelenleri* (*Galanthus* L.) I. Ege Üniversitesi Basımevi, Bornova, Izmir (1995).
99. Davis, A.P., Brickell, C.D., *Galanthus peshmenii*: a new snowdrop from the eastern Aegean. *The New Plantsman*, 1, 1, 14-19 (1994).
100. Kerndorff, H., Two new taxa in Turkish *Crocus* (*Iridaceae*). *Herbertia*, 49, 1-2, 76-86 (1993).
101. Kerndorff, H., Pasche, E., Zwei bemerkenswerte Taxa des *Crocus biflorus* - Komplexes (*Iridaceae*) aus der Nordosttürkei. *Linzer Biol. Beitr.* 29, 1, 591-600 (1997).
102. Mathew, B., An interesting new autumn - flowering *Crocus* from Turkey. *The Royal Horticultural Society, The New Plantsman*, 2, 3, 182-184 (1995).
103. Pasche, E., A new *Crocus* (*Iridaceae*) from Turkey. *Herbertia* 49, 1-2, 67-75 (1993).
104. Kerndorff, H., Pasche, E., *Crocus mathewii*. A new autumn-flowering *Crocus* from Turkey. *The New Plantsman*, 1, 2, 102-106 (1994).
105. Baytop, A., The presence of *Cyperus glomeratus* L. in Turkey. *Doğa Tu Botanik D.*, 20, 6, 535-36 (1996).
106. Scholz, H., Byfield, A., Three grasses new to Turkey. *Doğa Tu Botanik D.* (1998) –baskıda-
107. Soreng, R.J., Hein, P. & Scholz, H., *Poa akmanii* (*Poaceae*), a new species from Turkey. *Willdenowia*, 27, 195-198 (1997).
108. Yıldırımlı Ş., *Spergularia sezer-zenginii* Yıldırımlı (*Caryophyllaceae*). a new species from Turkey. *Ot Sistematisk Botanik Dergisi*, 4, 2, 1-6 (1997).
109. Erik, S. & Demirkuş, N., New species from NE Turkey: *Chaerophyllum posofianum* (*Apiaceae*) and *Vicia erzurumica* (*Fabaceae*). *Willdenowia* 28, 151-156 (1998).
110. Wagenitz, G., Ertuğrul, K. & Dural, H., A new species of *Centaurea* sect. *Psephelloideae* (*Compositae*) from SW Turkey. *Willdenowia* 28, 157-161 (1998).
111. Alpinar, K., Türkiye'de *Carduncellus caeruleus* (L.) C.Presl var.*incisus* DC.'un varlığı. *Ot Sistematisk Botanik Dergisi*, 4, 1, 43-48 (1997).